


**Claims**


1. A process for preparing a compound of formula (I):



(I)

or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, wherein:


- 10 T represents a substituted or unsubstituted aryl group and T<sup>1</sup> is O or S; which process comprises, treating a compound of formula (II):



(II)


or a tautomeric form thereof and/or a salt thereof and/or a solvate thereof, wherein T and T<sup>1</sup> are as defined in relation to formula (I), with a microbial reductase obtained from an appropriate red yeast; and thereafter, as required, preparing a pharmaceutically acceptable salt and/or a pharmaceutically acceptable solvate of the compound of formula (I) or a tautomeric form thereof.

2. A process according to claim 1, wherein T represents a moiety selected from the list consisting of (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ii), (Ij), (Il), (Im), (In), (Io) and (Ip):




(Ia)

wherein  $\text{A}^1$ ,  $\text{A}^2$ ,  $\text{R}^1$  and  $n$  are as defined in relation to formula (I) of EP  
 5 0306228;




(Ib)

wherein  $\text{R}^2$ ,  $\text{L}^1$ ,  $\text{L}^2$  and  $\text{L}^3$  are as defined in relation to formula (I) of  
 10 EP 0008203;



(Ic)

15 wherein  $\text{R}^1$ ,  $\text{R}^2$ ,  $\text{R}^3$ ,  $\text{R}^4$ ,  $\text{R}^5$ ,  $\text{W}$  and  $n$  are as defined in relation to formula  
 (I) of EP 0139421;




(Id)

20


wherein  $\text{R}^1$ ,  $\text{R}^2$  and  $\text{R}^3$  are as defined in relation to formula (I) of  
 EP 0032128;

-25-




(Ie)

wherein A, R, R<sup>1</sup> and X are as defined in relation to formula (I) of  
 5 EP 0428312;




(If)

10 when A, B, R and R<sup>1</sup> are as defined in relation to formula (II) of  
 EP 0428312;




(Ig)

15 wherein R<sup>1</sup> is as defined in relation to formula (I) of EP 0489663;



(Ih)


20 wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and n are as defined in relation to formula (I) of  
 EP 0155845;



(Ii)


when R<sup>1</sup> is as defined in relation to formula (I) of EP 0257781;

5




(Ij)

10 wherein Ar, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, n, U and W are as defined in relation to formula (I) of United States Patent No. 5104888;



(Ik)


15 when A, R<sup>1</sup>, R<sup>2</sup> and X are as defined in relation to formula (I) of EP 0208420;



(Il)

20


when R<sup>1</sup>, R<sup>2</sup>, X, Z m and n are as defined in relation to formula (I) of EP 0177353;



(Im)


according to formula (I) of EP 0319189;

5




(In)

10 wherein A, B, X, X<sup>1</sup>, X<sup>2</sup>, n and Z are as defined in relation to formula (I) of EP 0332331;



(Io)


15 wherein V, W, X, Y, Z, Z<sup>1</sup> and n are as defined in EP 0332332; and




(Ip)

20 wherein Q and X are as defined in relation to formula (I) of International Application No. WO 92/18501.


3. A process according to claim 1 or claim 2, wherein T represents a moiety selected from the list consisting of (a), (b), (c), (d), (e), (f), (g), (h)  
25 and (i):




(a)




(b)




(c)



(d)




(e)




(f)


5



(g)



(h), and




(i)

4. A process according to any one of claims 1 to 3, wherein T represents a moiety of formula (Ia).

5. A process according to any one of claims 1 to 4, wherein T<sup>1</sup> represents S.

6. A process according to claim 1 for the preparation of a compound of formula (1):




(1)

10 or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof, and/or a pharmaceutically acceptable solvate thereof, wherein:  
 A<sup>1</sup> represents a substituted or unsubstituted aromatic heterocyclyl group;  
 R<sup>1</sup> represents a hydrogen atom, an alkyl group, an acyl group, an aralkyl group, wherein the aryl moiety may be substituted or unsubstituted, or a  
 15 substituted or unsubstituted aryl group;  
 A<sup>2</sup> represents a benzene ring having in total up to five substituents; and  
 n represents an integer in the range of from 2 to 6;

which process comprises, treating a compound of formula (2):


20



(2)

or a tautomeric form thereof and/or a salt thereof, and/or a solvate thereof,  
 25 wherein A<sup>1</sup>, A<sup>2</sup>, R<sup>1</sup> and n are as defined in relation to formula (1) with a microbial reductase obtained from an appropriate red yeast; and thereafter, as required, preparing a pharmaceutically acceptable salt, and/or a pharmaceutically acceptable solvate of the compound of formula (1) or a tautomeric form thereof.

7. A process according to claim 6, wherein the compound of formula (1) is 5-(4-[2-N-methyl-N-(2-pyridyl)amino]ethoxy)benzyl)-2,4-thiazolidinedione, or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof, and/or a pharmaceutically acceptable solvate thereof.
8. A process according to any one of claims 1 to 7, wherein an appropriate red yeast is a red yeast which provides the above mentioned reduction, including known red yeasts and those red yeasts which may be produced from known red yeasts by conventional methods.
9. A process according to any one of claims 1 to 8, wherein an appropriate red yeast is a red yeast from the species of the genera *Rhodotorula*, *Rhodospiridium* or synonyms thereof.
10. A process according to any one of claims 1 to 9 wherein an appropriate red yeast is *Rhodotorula glutinis* CBS 4406, *Rhodotorula rubra* CBS 6469, *Rhodotorula rubra* CBS 17 and *Rhodotorula glutinis* IFO 0869.
11. A process according to any one of claims 1 to 9 wherein an appropriate red yeast is *Rhodospiridium toruloides* CBS 14
12. A process for the preparation of a compound of formula (I) (the 'enantiomerically enriched compound (I)') wherein greater than 50% w/w of said compound is in the form of a compound of formula (IA) :



(IA)

or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, wherein T and T<sup>1</sup> are as defined in relation to formula (I) and the \*\*\* carbon atom is an enantiomeric carbon atom, which process comprises reacting a



compound of the above defined formula (II) with a microbial reductase  
btained from an appropriate red yeast and wherein the reaction is  
carried out at an acidic pH; and thereafter, as required, preparing a  
pharmaceutically acceptable salt and/or a pharmaceutically acceptable  
5 solvate of the enantiomerically enriched compound (I) or a tautomeric  
form thereof.

13. Enantiomerically enriched compound (I) or a tautomeric form  
thereof and/or a pharmaceutically acceptable salt thereof and/or a  
10 pharmaceutically acceptable solvate thereof, wherein greater than 50%  
w/w is in the form of compound (IA)

14. A compound of formula (IA) or a tautomeric form thereof and/or a  
pharmaceutically acceptable salt thereof and/or a pharmaceutically  
15 acceptable solvate thereof, in optically pure form.

15. Enantiomerically enriched compound (I), or a tautomeric form  
thereof and/or a pharmaceutically acceptable salt thereof and/or a  
pharmaceutically acceptable solvate thereof, for use as an active  
20 therapeutic substance.

16. Enantiomerically enriched compound (I), or a tautomeric form  
thereof and/or a pharmaceutically acceptable salt thereof and/or a  
pharmaceutically acceptable solvate thereof, for use in the treatment of  
25 and/or prophylaxis of hyperglycaemia, hyperlipidaemia, hypertension,  
cardiovascular disease and certain eating disorders.

17. A pharmaceutical composition comprising enantiomerically  
enriched compound (I), or a tautomeric form thereof, or a  
30 pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable  
solvate thereof, and a pharmaceutically acceptable carrier therefor.